


# Basin Overview November 2003

# Lower Eastern Shore



## Restoring the Bay:

### The Land-Water Connection

The Lower Eastern Shore, like other tributaries to the Chesapeake Bay, is degraded by nutrient and sediment pollution harming aquatic life. Excess nutrients and sediments are the primary sources of pollution in the Chesapeake Bay. Nutrients occur naturally in soil, animal waste, plants, and the atmosphere; but in the Chesapeake Bay watershed, urbanization and farming have increased nutrient loads to unhealthy levels. These nutrients – nitrogen and phosphorus – promote the growth of algae, which in turn, blocks sunlight from reaching underwater grasses and reduces dissolved oxygen and suitable habitat for aquatic life.

### The Lower Eastern Shore Basin

The Lower Eastern Shore basin drains approximately 1,450 square miles of land in Wicomico and Somerset Counties and a portion of Caroline, Dorchester and Worcester Counties.

Major water bodies in the basin include the Pocomoke, Manokin, Big Annemessex, Nanticoke, Wicomico, Blackwater and Transquaking Rivers, Marshyhope and Monie Creeks, and embayments such as Tangier Sound, Fishing Bay and Honga River.

The Lower Eastern Shore, along with all tributary basins in the Chesapeake, contribute to and are impacted by nutrient pollution. Nutrient pollution can be divided into two major categories – point sources (pollution that comes from a single, definable location, such as a wastewater treatment plant or industrial discharge) and non-point sources (pollution that cannot be attributed to a clearly identifiable, specific physical location, such as runoff from land and atmospheric deposition). Runoff from different land uses, point sources, and atmospheric deposition are the major sources of nutrients within the Bay watershed.

In the Lower Eastern Shore basin, land use is

primarily agricultural and forests. Because of the agricultural nature of the Lower Eastern Shore, the largest portions of the nitrogen and phosphorus pollutants come from agriculture sources. The remaining contributions come from a combination of non-point and point sources.


In this basin, as in the rest of the state, forest and wetlands are a land use that releases few nutrients to rivers and the Bay. Baywide, approximately 33% of nitrogen loads come from atmospheric sources, however, that varies from basin to basin and is included in land based loads.

### A Work in Progress

Maryland has been working since the first Chesapeake Bay Agreement was signed in 1983 to reduce nutrient pollution to the Chesapeake Bay. Since 1985, wastewater treatment plants, farmers, and others have achieved significant nitrogen and phosphorus reductions. Nitrogen loads in the Lower Eastern Shore basin have been reduced 30% from 9.46 to 6.66 million pounds a year since 1985, and phosphorus loads have been reduced 52% from 1.10 to .53 million pounds.

Large portions of these reductions were achieved through agricultural best management practices (BMPs).

### Lower Eastern Shore Basin Nutrient Goals



\*Updated 2002 Progress information available soon

These practices provide the most effective and practicable means of controlling pollutants, such as nutrient management or cover crops. In the Lower Eastern Shore basin, nitrogen loads from agriculture dropped 43% and phosphorus decreased 63%.

## Goals for a Healthy Bay

In 2000, the Chesapeake Bay Program partners – Maryland, Virginia, Pennsylvania, the District of Columbia, the U.S. Environmental Protection Agency, and the Chesapeake Bay Commission – signed *Chesapeake 2000*, a new agreement designed to protect and restore living resources, vital habitats, and water quality in the Bay and its watershed. Key parts of this agreement include developing new nutrient and sediment goals for the Bay and its tidal tributaries based on the needs of living resources and revising the Tributary Strategies to achieve these new goals.

In the spring of 2003, the Chesapeake Bay Program finished developing water quality criteria that identify the levels of dissolved oxygen, water clarity, and chlorophyll (algae) which are needed to support healthy populations of Bay living resources. The Chesapeake Bay Program used computer models to estimate the amount of nitrogen and phosphorus loads (also called loading caps) that can enter the Bay while achieving these water quality criteria. These loads were allocated to each tributary basin and state. As a result, each basin will have nutrient reductions to be achieved in order to reach their nutrient loading cap.

*The revised nutrient caps for the Lower Eastern Shore basin are 3.78 million pounds of nitrogen and .33 million pounds of phosphorus.*




## Water Quality

The monitoring data for the Lower Eastern Shore shows trends of how water quality has changed over time from 1985 to 2001. Over this period, nitrogen levels are relatively good in the sounds but are poor in many areas, and have worsened at the Sharpstown station on the Nanticoke River over the 1985-2001 period. Phosphorus levels are relatively fair to good, and have improved in some areas. However, algae levels have increased in Fishing Bay and Tangier Sound. Total suspended solid levels are poor throughout most of the basin. Although there are no trends over the 1985-2001 period, TSS levels appear to have been improving in the most recent years.


## Living Resources in the Lower Eastern Shore

The basin supports more than 100 species of fish in its freshwater streams and brackish waters, including herring, white and yellow perch, bluefish and flounder. This area also supports an important commercial and recreational blue crab

## Total Nitrogen Concentrations: Lower Eastern Shore



## Total Phosphorus Concentrations: Lower Eastern Shore



fishery and serves as a nesting and overwintering area for bald eagles.

### Local Benefits

By addressing nutrient and sediment pollution in the Lower Eastern Shore basin there will be local and downstream advantages. The overall result will be a decrease in algal production which will aid in the return of underwater grasses and improved habitat.

For nontidal areas, the Maryland Biological Stream Survey (MBSS) provides a picture of overall ecological stream health (since 1995 in this basin). Data, such as measures of the variety of species, pollution sensitivity, and proportion of exotic species, are collected for each stream. These data are combined into one overall value, or index of health, for the streams in the Lower Eastern Shore watershed that is referred to as an Index of Biotic Integrity (IBI). By using this index, complex ecological information can be summarized and stream health can be rated as good, fair, poor, or very poor. Streams rated good or fair by the index are considered healthy compared to reference streams, while streams rated poor or very poor are considered unhealthy.

In the Nanticoke and Pocomoke Rivers a large portion of the of the monitoring sites were rated as having a fair Index of Biotic Integrity scores. The Nanticoke has the highest percentages of wetlands of any stream basin in the state. The Pocomoke River basin contains 2,700 acres of land considered to be of exceptional ecological importance. The basin has been described as having one of the best environments for bird life on the Atlantic Coast.

Addressing the quality of the streams will translate into local habitat quality and contribute to the support of such critical natural resources to the Bay. Healthy local streams and rivers will not just serve as a recreational asset to the local community but often translates into an increased quality of life and local economic benefits

### Downstream Benefits

Restoration efforts in the Lower Eastern Shore will be felt elsewhere. By achieving our nutrient goals, and addressing sediment in the Lower Eastern Shore basin, we expect to see decreased algal production downstream, better habitat and a resurgence of underwater grasses. The following is a description of living resource challenges and goals for the mainstem and tidal areas of the Chesapeake Bay Watershed.

### Bay Grasses

Underwater grasses, or submerged aquatic vegetation, play an important ecological role to the Chesapeake Bay environment. They provide food, refuge and nursery habitat for many waterfowl, fish, shellfish and invertebrates, and



produce oxygen in the water column. These grasses also filter and trap sediment that cloud the water and bury bottom-dwelling organisms, such as oysters; provide shoreline erosion protection by slowing down wave action; and remove excess nutrients that could fuel unwanted growth of algae in the surrounding waters.

Submerged aquatic vegetation had largely vanished in the Bay by the 1970s, primarily due to poor water quality. Over the past decade, improvements in water quality have led to a modest resurgence in underwater grasses in some parts of the Bay. In 2000, underwater grasses covered about 69,000 acres in the Bay. In 2003, the Chesapeake Bay Program set a new goal for underwater grasses of 185,000 acres Baywide. This was based in part on the amount of grasses that would return once we achieve the new nutrient reduction goal.

### Blue Crabs

The blue crab is one of the most important species harvested in the Bay. It has the highest value of any commercial fishery and supports a recreational fishery of significant, but undetermined, value. Due to loss of habitat and harvest pressure, however, the abundance of mature female crabs is at near historic lows. The *Chesapeake 2000 Agreement* calls for the Bay partners to “manage the blue crab fishery to restore a healthy spawning biomass, size, and age structure.” To achieve this, Maryland and Virginia have committed to reduce harvest pressure on blue crabs by 15% compared to the harvests of 1997 through 1999.

Restoring underwater grasses will be an important step in restoring blue crab populations. During the 1970s and 1980s, the widespread disappearance of underwater grasses resulted in a severe loss of important crab habitat and nursery areas, primarily for females and crabs in the molting stage. Bay scientists have found that 30 times more juvenile crabs were found in areas with Bay grasses than in areas without.



## Oysters

Over-harvesting, dwindling habitat, pollution, and diseases (such as Dermo and MSX) have caused a severe decline in oysters throughout the Chesapeake Bay over the last century. Since the 1950s, harvests have fallen Baywide from 35 million pounds to below 3 million pounds. In addition to their fisheries value, oysters are critical to the Bay's ecosystem. They provide habitat for many Bay species and help improve water clarity by filtering algae and sediment from the water.

The *Chesapeake 2000 Agreement* commits to increasing native oysters tenfold by 2010. The Oyster Restoration Strategy, which was developed to support the agreement, focuses on rehabilitating oyster habitat, much of which is degraded by silt and nearly barren. In addition to improving habitat, the strategy aims to increase the oyster population by the construction of a Baywide network of non-harvest sanctuary areas. Up to 250 such areas have been suggested throughout the Bay so far. Protected from harvesting, it is hoped that some of the oysters in these sanctuaries will survive disease and enhance the Bay's oyster population.



by the team and the Tributary Strategies Development Workgroup. This workgroup worked closely with state and local governments, team members, local constituents and other stakeholder groups. The working draft of the document will be available for review, and a second round of public meetings is planned for December 2003 for public review of the strategies.

## Tools for Change

Maryland's Tributary Teams are leading the revision of their Tributary Strategies – watershed-based plans to achieve the nutrient and sediment goals within each of the state's 10 tributary basins. Restoring the Lower Eastern Shore will require the active involvement of all watershed residents. Strategies for the Lower Eastern Shore basin will be drawn from an array of measures to reduce the amounts of nutrients from wastewater treatment plants and agricultural, urban, and suburban lands. Protection of forests and wetlands will help prevent increases in nitrogen and phosphorus loads.

## The Next Steps

Over the coming months, the Lower Eastern Shore Tributary Team and Maryland's Departments of Natural Resources, Environment, Agriculture, and Planning will work closely with residents of the basin to identify best management practices that can be applied in the watershed to reduce nutrient pollution and restore habitat.

These practices will be summarized in a Tributary Strategy for the basin. Funds to implement this strategy will be sought from federal, state, and local governments. Private landowners and other watershed residents will also contribute. While implementation may not be complete by the target date of 2010, every effort will be made to reach the water quality goals by that date.

With input from the 1st public meeting in June 2003, the Tributary Strategy for the Lower Eastern Shore basin was drafted this summer and fall

### For more information or how to get involved with the Lower Eastern Shore Tributary Team:

- Lower Eastern Shore Technical Basin Summary: [www.dnr.state.md.us/bay/tribstrat/basin\\_summaries.html](http://www.dnr.state.md.us/bay/tribstrat/basin_summaries.html)
- Maryland Biological Stream Survey: Pocomoke: [www.dnr.state.md.us/streams/pubs/pocomoke.pdf](http://www.dnr.state.md.us/streams/pubs/pocomoke.pdf)
- Nanticoke: [www.dnr.state.md.us/streams/pubs/nanticoke.pdf](http://www.dnr.state.md.us/streams/pubs/nanticoke.pdf)
- Chesapeake Bay water quality criteria: [www.chesapeakebay.net](http://www.chesapeakebay.net)
- Maryland's water quality standards: [www.mde.state.md.us](http://www.mde.state.md.us)
- Maryland's Tributary Teams: [www.dnr.state.md.us/bay/tribstrat.html](http://www.dnr.state.md.us/bay/tribstrat.html)

Look for the next round of Tributary Strategy public meetings or get involved with your local Tributary Team!

Lower Eastern Shore Team Coordinator, Stephanie Blades, at [sblades@dnr.state.md.us](mailto:sblades@dnr.state.md.us) or 410.260.8723

Produced by the MD Department of Natural Resources with funds from the Environmental Protection Agency, Chesapeake Bay Program. This document does not necessarily reflect the opinion of the EPA. Printed on recycled paper.